arXiv:hep-ph/9702259v1 5 Feb 1997

Lower bound for the ground state energy of the no-pair Hamiltonian

C. Tix*
Matematisk institutt, Universitetet i Oslo
Postboks 1053, N-0316 Oslo, Norway

PACS: 03.65.Pm, 03.65.Ge, MCS: 35P15, 35Q40†

14 January 1997

Abstract

A lower bound for the ground state energy of a one particle relativistic Hamiltonian - sometimes called no-pair operator - is provided.

1 Introduction

The multi-particle analogon of the operator

$$B = \Lambda_+ \left(D_0 - \frac{e^2 Z}{|\boldsymbol{x}|} \right) \Lambda_+ \tag{1}$$

is often used to describe relativistic effects in atoms and molecules, see, e.g., [8, 9]. Here D_0 is the free Dirac operator $D_0 = c\frac{\hbar}{i}\boldsymbol{\alpha}\cdot\nabla + \boldsymbol{\beta}mc^2$ and

$$\Lambda_+(\boldsymbol{p}) = \frac{1}{2}\left(1 + \frac{c\boldsymbol{\alpha}\cdot\boldsymbol{p} + \boldsymbol{\beta}mc^2}{E(\boldsymbol{p})}\right)$$

with $E(\boldsymbol{p}) = (c^2\boldsymbol{p}^2 + m^2c^4)^{1/2}$ is the projection operator on the electron subspace of the free Dirac operator in momentum space. The underlying Hilbert space of B is the space of square integrable four spinors ψ with $\Lambda_+\psi = \psi$.

The operator B was introduced for the two particle case by Brown and Ravenhall [2] (see also [1, 14]) to cure the continuum dissolution of the two particle Dirac operator with Coulomb interaction: since the Dirac operator is unbounded form below the two particle Dirac operator has the whole real line as spectrum, the eigenvalues of the one particle operator “dissolve”. B is called

---

[*] E-mail: christti@math.uio.no

[†] Keywords: relativistic, wave equation, bound, ground state

no-pair operator or in the description of meson models (usually with different potentials) "reduced Salpeter operator" [11]. Recently, its multi particle analogon was used in connection with the "stability of relativistic matter"–question [10].

Similar to the Dirac operator with Coulomb potential, the no-pair operator has a critical nuclear charge Z_c. Hardekopf and Sucher [6, 7] observed that $Z_c = 2/[(\pi/2+2/\pi)\alpha]$ with the fine structure constant $\alpha := e^2/(\hbar c)$ and investigated the ground state energy of B numerically. They claimed that, as for the Dirac equation, the ground state energy vanishes for $Z = Z_c$ [7, p. 2025]. Evans et al [3] proved that the energy $(\psi, B\psi)$ is bounded from below by $\alpha Z(1/\pi - \pi/4)mc^2$, if the nuclear charge does not exceed the critical charge Z_c, otherwise it is unbounded. Tix [15] improved this bound to $mc^2\,(1 - \alpha Z - 0.002Z/Z_c) \geq 0.09$. This last result shows the difficulties to obtain accurate numerically results near the critical coupling parameter and the need for sharp bounds. In this short note we explore the results from [15] by using some rather trivial numerics to obtain a reliable lower bound for the ground state energy for all $Z \leq Z_c$.

2 The Lower bound

A short summary of the method to obtain a lower bound for $(\psi, B\psi)$ given in [3, 15] is provided. Any normalized four spinor in the electron subspace of D_0 can in be written in momentum space as

$$\widehat{\psi}(\boldsymbol{p}) = \frac{1}{N(\boldsymbol{p})} \begin{pmatrix} (E_0 + E(\boldsymbol{p}))\, u(\boldsymbol{p}) \\ c\boldsymbol{p}\cdot\boldsymbol{\sigma} u(\boldsymbol{p}) \end{pmatrix} \tag{2}$$

with the normalized Pauli spinor u, the Pauli-matrices $\boldsymbol{\sigma}_j$ and $E_0 = E(\boldsymbol{0})$, $N(\boldsymbol{p}) = [2E(\boldsymbol{p})(E(\boldsymbol{p}) + E_0)]^{1/2}$. In [3] it was shown that the minimizer of $(\psi, B\psi)$ is radially symmetric, i.e., the minimizing u is of the form $u(\boldsymbol{p}) = a(|\boldsymbol{p}|)(\sqrt{4\pi}|\boldsymbol{p}|)^{-1}(1,0)^*$ where a is normalized and positive. This yields for the energy $(\psi, B\psi)$

$$\int_0^\infty e(p)a(p)^2 dp - \frac{\alpha Z}{2\pi}\int_0^\infty \int_0^\infty a(p')k(p',p)a(p)dpdp' \tag{3}$$

with $e(p) = E(\boldsymbol{p})$, $p = |\boldsymbol{p}|$, $k := k_0 + k_1$ and

$$k_j(p', p) := s_j(p')\, Q_j\left(\frac{p}{2p'} + \frac{p'}{2p}\right) s_j(p). \tag{4}$$

The functions Q_l are Legendre functions of the second kind (see [13] for the notation) and appear here for the same reason as in the treatment of the non–relativistic hydrogen atom in momentum space [4, problem 77] and $s_j(p) = \sqrt{1 + (-1)^j/e(p)}$. Because of scaling we have assumed $\hbar = m = c = 1$ in (3). Using the Schwarz inequality to introduce trial functions $h_j(p) > 0$ in the potential energy part of $(\psi, B\psi)$

$$\begin{aligned}\int_0^\infty \int_0^\infty & a(p')k_j(p',p)a(p)dpdp' \\ &= \int_0^\infty \int_0^\infty a(p')s_j(p')\sqrt{Q_j}\,\sqrt{\frac{h_j(p)}{h_j(p')}}\,\sqrt{\frac{h_j(p')}{h_j(p)}}\,\sqrt{Q_j}\;a(p)s_j(p)\,dpdp' \\ &\qquad\qquad \le \int_0^\infty a(p)^2 s_j(p)^2 \int_0^\infty \frac{h_j(p')}{h_j(p)} Q_j \left(\frac{p}{2p'} + \frac{p'}{2p}\right) dp'\,dp\end{aligned}$$

is obtained. Note that the operator B is similar to the Herbst operator $H = \sqrt{c^2p^2 + m^2c^4} - e^2Z/|\boldsymbol{x}|$. The ground state energy of H is obtained by minimizing the quadratic form (3) with $k = k_0$, without the square roots and with $2Z$ instead of Z. Since no error is introduced by using the Schwarz inequality when h_j/s_j is proportional to the ground state wave function of (3), this suggests a similar trial function than the one that was used by Raynal et al [12] for the Herbst operator. They obtained excellent upper and lower bounds for the ground state of H. Choosing the trial functions h_j essentially as the Fourier transform of $|\boldsymbol{x}|^{\beta-1}e^{-\mu|\boldsymbol{x}|}$, the lower bound

$$(\psi, B\psi) \ge \sup_{\substack{-1<\beta<1\\ \mu>0}} \inf_{p>0} \mathcal{E}[Z,\beta,\mu](p). \tag{5}$$

was obtained in [15] where

$$\begin{aligned}\mathcal{E}[Z,\beta,\mu](p) = e(p) &- \frac{\alpha Z}{2}\frac{\sqrt{p^2+\mu^2}}{\beta}\frac{\sin[\beta\arctan(p/\mu)]}{\sin[(\beta+1)\arctan(p/\mu)]}\left(1 + \frac{1}{e(p)}\right) \\ &- \frac{\alpha Z}{2}\left(1 - \frac{1}{e(p)}\right)\frac{\sqrt{p^2+\mu^2}}{\beta+2}\,\frac{P^{-3/2}_{\beta-1/2}\left(\mu(p^2+\mu^2)^{-1/2}\right)}{P^{-3/2}_{\beta+1/2}\left(\mu(p^2+\mu^2)^{-1/2}\right)}\end{aligned} \tag{6}$$

with the associated Legendre functions $P_\nu^{-3/2}$ (see [5, p. 1060]). To cancel the large momentum terms in (6) $\beta \ge 0$ is chosen to fulfill

$$\frac{2}{\alpha Z} = \frac{\tan(\beta\pi/2)}{\beta} + \frac{\beta}{(1-\beta^2)\tan(\pi\beta/2)}. \tag{7}$$

The resulting function $\mathcal{E}[Z,\beta,\mu]$ is numerically evaluated. The values for the parameters β and μ for some values of αZ can be found in table 1 and a plot of the energy over the coupling constant is shown in figure 1.

αZ	E	μ	β
0.9060	0.106	0.9843	0
0.9003	0.215	0.9526	0.1
0.8527	0.423	0.8652	0.3
0.7500	0.621	0.7395	0.5
0.5709	0.808	0.5552	0.7

Table 1: Lower bound for the energy $E = \inf_p \mathcal{E}[Z, \beta, \mu](p)$ and the corresponding parameters



Figure 1: Lower bound for the energy $(\psi, B\psi)$ of the no-pair Hamiltonian

3 Acknowledgment

This work was partially supported by the European Union under grant ERB4001GT950214 and under the TMR-network grant FMRX-CT 96-0001.

References

[1] Hans A. Bethe and Edwin E. Salpeter. Quantum mechanics of one- and two-electron atoms. In S. Flügge, editor, *Handbuch der Physik, XXXV*, pages 88–436. Springer, Berlin, 1 edition, 1957.

[2] G.E. Brown and D.G. Ravenhall. *Proc. Roy. Soc. London A*, 208(A 1095):552–559, September 1951.

[3] William Desmond Evans, Peter Perry, and Heinz Siedentop. *Commun. Math. Phys.*, 178(3):733–746, July 1996.

[4] Siegfried Flügge. *Practical Quantum Mechanics I*, volume 177 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1 edition, 1982.

[5] I.S. Gradshteyn and I.M. Ryzhik. *Table of integrals, series, and products.* Academic Press, New York and London, 4 edition, 1965.

[6] G. Hardekopf and J. Sucher. *Phys. Rev. A*, 30(2):703–711, August 1984.

[7] G. Hardekopf and J. Sucher. *Phys. Rev. A*, 31(4):2020–2029, April 1985.

[8] Y. Ishikawa and K. Koc. *Phys. Rev. A*, 50(6):4733–4742, December 1994.

[9] Hans Jörgen Aa. Jensen, Kenneth G. Dyall, Trond Saue, and Knut Fægri Jr. *J. Chem. Physics*, 104(11):4083–4097, March 1996.

[10] Elliott H. Lieb, Heinz Siedentop, and Jan Philip Solovej. Stability and instability of relativistic electrons in classical electromagnetic fields. *J. Stat. Phys.*, submitted, 1997.

[11] M. G. Olsson, S. Veseli, and K. Williams. *Phys. Rev. D*, 53(1):504–513, 1996.

[12] J. C. Raynal, S.M. Roy, V. Singh, A. Martin, and J. Stubbe. *Phys. Lett. B*, 320:105–109, 1994.

[13] Irene A. Stegun. Legendre functions. In Milton Abramowitz and Irene A. Stegun, editors, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, chapter 8, pages 331–353. Dover Publications, New York, 1965.

[14] J. Sucher. *Phys. Scripta*, 36:271–281, 1987.

[15] C. Tix. Strict positivity of a relativistic hamiltonian due to Brown and Ravenhall. *Bull. London Math. Soc.*, submitted, 1997.